

Seed priming induces biochemical changes in melon plants and increase salt tolerance

CASTAÑARES, J.L.^{1,2}; BOUZO, C.A.³

ABSTRACT

An experiment was conducted to evaluate the effect of priming on germination and initial growth of melon plants under salinity conditions. Two osmotic agents (NaCl and CaCl₂) and two treatment durations (2 and 4 days) were compared. Germination percentage and growth of seedlings were evaluated at electrical conductivity 8.0 dS m⁻¹. Total chlorophyll content, relative water content, root viability, proline content, relative electrolyte leakage, malondialdehyde concentration, peroxidase and catalase activity, Na⁺/K⁺ ratio in green parts, and K⁺ leakage from roots were measured. The best germination was obtained when seeds were soaked either with NaCl or CaCl₂ for two days. Priming improved growth parameters and plants presented a better response in all the evaluated biochemical parameters. Results suggest that priming could be used to improve the performance of seeds and seedlings in situations of high salinity.

Keywords: *Cucumis melo*, osmo priming, germination, salt stress, oxidative stress.

RESUMEN

Se realizó un experimento para evaluar el efecto del 'priming' de semillas sobre la germinación y crecimiento inicial de plantas de melón en salinidad. Se compararon dos agentes osmóticos (NaCl y CaCl₂) y dos duraciones de tratamiento (2 y 4 días). Se evaluó el porcentaje de germinación y crecimiento de las plántulas a una conductividad eléctrica de 8.0 dS m⁻¹. Se midió el contenido total de clorofila, contenido relativo de agua, viabilidad de raíces, contenido de prolina, pérdida relativa de electrolitos, concentración de malondialdehído, actividad peroxidasa y la catalasa, proporción de Na⁺/K⁺ en partes verdes y pérdida de K⁺ por raíces. La mejor germinación se obtuvo cuando las semillas se trataron con NaCl o CaCl₂ durante dos días. El 'priming' mejoró los parámetros de crecimiento y las plantas mostraron una mejor respuesta en todos los parámetros bioquímicos evaluados. Los resultados sugieren que el 'priming' podría usarse para mejorar el rendimiento de las semillas y las plántulas en situaciones de alta salinidad.

Palabras clave: *Cucumis melo*, osmo priming, germinación, estrés salino, estrés oxidativo.

¹Instituto Nacional de Tecnología Agropecuaria (INTA), Estación Experimental Agropecuaria (EEA) Área Metropolitana Buenos Aires (AMBA). Udaondo 1695, Ituzaingó, Buenos Aires, Argentina. Correo electrónico: castanares.jose@inta.gob.ar

²Universidad Nacional de Luján (UNLu), Departamento de Ciencias Básicas, Laboratorio de Fisiología Vegetal. Ruta 5 y Avenida Constitución, Luján, Buenos Aires, Argentina.

³Universidad Nacional del Litoral, Facultad de Ciencias Agrarias, Laboratorio de Biología Molecular y Fisiología Vegetal. Kreder 2805, Esperanza, Santa Fe, Argentina.

INTRODUCTION

Soil salinity is one of the most important stresses limiting crop production. This phenomenon affects 20% of cultivated lands of the world and causes important yield losses in crops (Qadir *et al.*, 2014). Irrigation with poor quality water is the main cause of increased soil salinity in cultivated land (Yadav *et al.*, 2011). Although irrigated crops represent only 15% of total cultivated land, this represents one third of the world's food production given that yields double what is produced in non-irrigated crops (Munns, 2005). For this reason, it is unlikely that crop irrigation will be abandoned as a normal practice and, instead, an increase in its application could be expected (Shabala and Munns, 2012).

Melon (*Cucumis melo* L.) is an important horticultural crop with a worldwide production of 27.3 million tons. China, Iran, Turkey, Egypt, and the United States represent 68% of the world's production (Iqbal *et al.*, 2016). It is cultivated in all regions of the world with warm and slightly rainy weather, where salinity is becoming a problem (Yasar *et al.*, 2006). This crop is considered as moderately sensitive to salt (salinity threshold in the range of 0.75 - 6.8 dS m⁻¹) (Tedeschi *et al.*, 2011). It has been reported that the highest sensibility occurs in early vegetative stages, while its tolerance increases between fruit development and harvest (Franco *et al.*, 1993; Nukaya *et al.*, 1984). High salinity can delay or inhibit germination (Carvajal *et al.*, 1998; Tanveer *et al.*, 2012), inhibit growth (Bokochoy-Nbok and Bokochoy, 1991; Botia *et al.*, 1998) and reduce yield (Huang *et al.*, 2012; Sivritepe *et al.*, 2005).

Seed priming is a technique of controlled hydration that allows activation of numerous metabolic processes (pre-germinative metabolism) which occur in early stages of imbibition, before radicle protrusion (Paparella *et al.*, 2015). There are different priming methods, which vary based on the way the entry of water is controlled: osmo-priming (seeds are soaked in an osmotic solution such as polyethylene glycol, mannitol or salts), hydropriming (seeds are soaked in a determined amount of water, or limited imbibition periods), or matrix priming (organic or inorganic solids are mixed with water to reduce osmotic potential) (Wojtyla *et al.*, 2016). Priming can improve seed germination under favorable and unfavorable conditions, although its effects are more evident under adverse conditions (Jisha *et al.*, 2013). The advantages of priming are not exclusive to the seed stage but it can also improve growth and yield of crops under different abiotic stressors (Castañares and Bouzo, 2017; Zheng *et al.*, 2016; Farhoudi and Sharifzadeh, 2006).

The objective of this work was to evaluate the effect of priming on germination, initial growth, and biochemical responses of melon plants grown in saline conditions.

MATERIALS AND METHODS

Seeds of F1 hybrid 'Charentais' cantaloupe 'Planter's Jumbo' (Bonanza Seeds, California, USA) were used. The

different reagents and salts (analytical grade) used were obtained from Sigma-Aldrich (Buenos Aires, Argentina).

Germination tests

Seeds were surface-disinfected using a 10% sodium hypochlorite solution (NaOCl in water, v/v) for 10 min. Then, the seeds were rinsed with deionized water, placed between two filter papers (Whatman N° 1) in 8.5 cm diameter Petri dishes, and soaked with either one of two different osmotic solutions: NaCl or CaCl₂, according to a previous study (Castañares and Bouzo, 2018), in a ratio of 1:5 (wt/v) (Farooq *et al.*, 2013). The concentration of these salts was calculated to achieve an osmotic potential (Ψ_o) of -1.5 MPa according to the van't Hoff equation (Shabala and Munns, 2012). Two priming durations were studied: two and four days. Ten seeds per dish were used, equidistant from each other, with five replicates per treatment. The assay was performed in a germination chamber (Semedic, model I-500 PF, Argentina) at 25 °C (\pm 0.5 °C) in darkness (Nascimento, 2003). After priming, seeds were removed, rinsed three times in distilled water, and air-dried at 25 °C for 24 h to reduce the moisture content to < 10%. Later, the seeds were germinated between two filter papers and imbibed with an 80 mM NaCl solution (EC = 8.0 dS m⁻¹). Control treatment (Control S1) consisted of seeds without priming, treated with distilled water. Germination was recorded daily during 8 days (ISTA, 2013) and seeds were labeled as germinated when the seed coat was broken and the radicle was visible (>2 mm). Germination percentage (GP) was calculated as the relation between germinated seeds and seeds put to germinate.

Growth of seedlings

The two treatments with the highest GP from the previous experiment were chosen to evaluate the growth response of seedlings. After priming, seeds were rinsed three times in distilled water and sown in 1000 cm³ pots previously filled with perlite, and watered with Hoagland's nutrient solution (Hoagland and Arnon, 1950), with 2.0 dS m⁻¹ electrical conductivity (EC). After first leaf emergence, plants started to be watered with the nutrient solution with the addition of approximately 60 mM NaCl, to achieve 8.0 dS m⁻¹ EC. EC was monitored daily with a portable conductivity meter (Milwaukee model WP MC66 0-10MS/C, USA). Two control groups were established: Control S0, plants derived from non-primed seeds and watered with a 2.0 dS m⁻¹ EC solution; and Control S1, plants derived from non-primed seeds and watered with an 8.0 dS m⁻¹ EC solution. The experiment was performed in a phytotron, with a 16 h light/8 h darkness cycle and temperatures of 25/18 °C (\pm 1.0 °C), optimal for melon vegetative growth (Reche Mármol, 2008).

The experiment had a completely randomized design. The position of the pots was randomly changed daily to minimize positional effects in the phytotron. All values reported are the means of five replicates. Data obtained were

analyzed using Tukey's test at 0.05 confidence level using Infostat statistical software (Di Rienzo *et al.*, 2011).

Growth evaluation

Five plants were extracted from each treatment after 40 days, and the following parameters were determined: leaf number, main stem height, leaf area, and total dry weight. Leaf area was determined using digitized images of leaves with Image J software (Schneider *et al.*, 2012). For stem height, the distance from the substrate surface up to the insertion of the petiole of the uppermost leaf was considered.

Xylematic potential measurement

Xylematic potential (Ψ_x) of plants was measured using a Scholander pressure chamber (Biocontrol model 0-6 MPA, Argentina). Measurements were made on developed leaves, located in the middle third of the stem (Scholander *et al.*, 1964), after 20 days of growth.

Determination of chlorophyll content, relative water content, root viability, proline and malondialdehyde concentration, and relative electrolyte leakage

After 20 days of growth in salinity, five plants per treatment were harvested and used for biochemical determinations.

Chlorophyll content (Ch) was determined according to Lichtenthaler and Wellburn (1983). A sample of 100 mg of fresh leaves (PF) was homogenized with 20 mL (V) of 80% (v/v) acetone in a mortar. Then the mixture was centrifuged at $8,000 \times g$ for 10 min. The absorbance (A) of supernatant was measured using a spectrophotometer (Shimadzu model UV-1800, Japan) at 646.8 and 663.2 nm, and chlorophyll content was determined according to the following formula:
$$\text{Ch (mg g}^{-1} \text{ FW)} = (7.15 A_{663.2 \text{ nm}} + 18.71 A_{646.8 \text{ nm}}) \times (V/\text{PF}).$$

Relative water content (RWC) was calculated as described by Cao *et al.* (2015). The second leaves were extracted and fresh weight (FW) was determined. Then the leaves were placed in a beaker with distilled water for 5 h to determine saturated weight (SW). Finally, the leaves were dried at 80 °C until constant weight and dry weight (DW) was determined. RWC was calculated according to the equation:
$$\text{RWC (\%)} = (\text{FW} - \text{DW}) / (\text{SW} - \text{DW}) \times 100.$$

Root viability was estimated by measuring the activity of the dehydrogenase enzyme by using the 2,3,5- triphenyl tetrazolium chloride (TTC) reduction technique (Clemensson-Lindell, 1994). A sample of 500 mg of fresh roots was cut into small pieces, put into test tubes with 5 mL of TTC 0.4 % and 5 mL of phosphate buffer (pH 7.0), and incubated for 3 h at 37 °C. The samples were then extracted in ethyl acetate for 15 min. The absorbance of samples was measured at 485 nm. Results were expressed as absorbance in relation to root dry weight, determined after drying in an 80 °C oven until constant weight ($A_{485 \text{ nm}} \text{ g}^{-1} \text{ DW}$).

Proline content was measured using the method developed by Bates *et al.* (1973). Approximately 300 mg of fresh

leaf material was homogenized in 10 mL of 3% aqueous 5-sulfosalicylic acid solution, and the homogenate was filtered using Whatman No 1 paper. An aliquot of 2 mL of filtrate was reacted with 2 mL of acid ninhydrin (1,2,3-triketohydrindene hydrate) and 2 mL of glacial acetic acid ($\text{C}_2\text{H}_4\text{O}_2$) for 1 h at 100 °C. The reaction was terminated on ice for 15 min. The reaction mixture was then extracted with 4 mL of toluene (C_7H_8) and vortexed for 20 s. The absorption of the upper phase was measured at 520 nm using a spectrophotometer and proline concentration was calculated using a standard curve and expressed as $\mu\text{mol g}^{-1} \text{ FW}$.

Relative electrolyte leakage (EL) was estimated according to Dionisio-Sese and Tobita (1998). A total of 100 mg of fresh leaves were cut into 5 mm segments and placed in test tubes with 10 mL of deionized water. The tubes were incubated 2 h at 30 °C and the initial EC of the medium was measured (EC1) using a conductivity meter (Altronix model CTX-II, USA). Then the test tubes were incubated at 100 °C for 15 min and the EC was measured again (EC2). EL was estimated according to the formula:
$$\text{EL (\%)} = \text{EC1/EC2} \times 100.$$

To evaluate the level of malondialdehyde (MDA), a sample of 300 mg of fresh leaves was macerated with 3 mL of 0.1% trichloroacetic acid ($\text{Cl}_3\text{C}_2\text{O}_2\text{H}$) (TCA). The homogenate was centrifuged at $3,000 \times g$ for 10 min. Then, 1 mL of the supernatant was placed in a test tube, and 1 mL of the reagent TCA-BHT-TBA (TCA 20%, thiobarbituric acid (TBA) 0.37% and butyl hydroxyl toluene (BHT) 0.01 g) was added. The sample was incubated in a water bath for 25 min at 95 °C. The reaction was terminated on ice. Then the sample was centrifuged at $10,000 \times g$ for 10 min. The supernatant was used to measure the absorbance at 532 and 600 nm. The content of MDA was expressed as $\mu\text{M g}^{-1} \text{ FW}$ (Heath and Packer, 1968).

Peroxidase and catalase activity

After 20 days of growth, fresh leaf samples (60 mg) were ground in a mortar, re-suspended with 1.5 mL of 10 mM phosphate buffer (pH 6), and centrifuged at $12,000 \times g$ for 20 min. The supernatant was used for the following assays.

To determine the peroxidase activity (POX), 100 μL of 10 mM sodium phosphate buffer (pH 6), 20 μL of guaiacol ($(\text{CH}_3\text{O})\text{C}_6\text{H}_4\text{OH}$) 0.25% (v/v) and 50 μL of 0.88 M H_2O_2 were added to 100 μL of the extract. A reduction in absorbance at 470 nm was measured at 0.5, 1.0, 1.5, and 2.0 min. POX activity was expressed as $\text{Abs min}^{-1} \text{ mg}^{-1} \text{ FW}$ (George, 1953).

Catalase activity (CAT) was measured according to Aebi (1984). An aliquot of 50 μL was extracted, and 2000 μL of 50 mM sodium potassium phosphate buffer (pH 7) and 20 μL of 0.88 M H_2O_2 were added. The activity was determined by measuring the decrease in absorbance due to H_2O_2 decomposition for 1.0 min at 240 nm and expressed as $\mu\text{mol of } \text{H}_2\text{O}_2 \text{ reduced g}^{-1} \text{ FW}$.

Determination Na^+/K^+ Ratio and K^+ leakage from roots

The following determinations were also made in seedlings after having grown for 20 days in salinity.

A sample of 2 g plant material (leaves and stems) was ground (2 mm) and placed in a muffle at 600 °C for 4 h. The ashes were boiled with 40 mL of HCl (30%) and 5 mL of HNO₃. The mixture was filtered and cooled. Deionized water was added to complete 250 mL (Campbell and Plank, 1998). K⁺ and Na⁺ contents were determined using a flame photometer (Metrolab, model 315, Argentina).

K⁺ leakage from roots was estimated according to Chen *et al.* (2005), in plants growing. Seedlings of each treatment were randomly chosen, their roots were washed and immersed in a beaker with 10 mL of 80 mM NaCl solution for 2 h. Then seedlings were removed, roots were surface dried with paper towels and root fresh weight was measured. The amount of K⁺ released into solution was determined using a flame photometer and expressed in mmol K⁺ g⁻¹ FW.

RESULTS

Germination in salinity

Priming with NaCl or CaCl₂ for two days reversed the inhibitory effect of salinity on melon seeds germination, compared to seeds without treatment (Control S1). Increasing treatment duration from 2 to 4 days reduced the favorable effect of the treatments of priming with NaCl and CaCl₂ (table 1).

Effect of priming on the growth of seedlings

Since the highest GP was registered for seeds exposed for two days with either salt, NaCl or CaCl₂, the fo-

Treatment	GP (%)
Control S1	72.00 ± 7.07 b
NaCl/2-d	98.00 ± 2.00 a
NaCl/4-d	68.00 ± 7.66 b
CaCl ₂ /2-d	94.00 ± 6.30 a
CaCl ₂ /4-d	74.00 ± 7.00 b

Table 1. Effect of priming with NaCl and CaCl₂ for 2 or 4 days on melon germination percentage (GP) at 8.0 dS m⁻¹. Different letters indicate a significant difference ($P \leq 0.05$) according to Tukey's test.

llowing determinations were made on plants using these treatments.

Priming allowed improving plant performance in salinity when compared to plants without treatment (fig. 1 and 2). In this way, priming allowed to increase the vegetative growth of plants measured through leaf number, main stem height, leaf area, and total dry weight.

Effect of priming on xylematic potential

Saline stress led to a reduction in xylematic potential (Ψ_x). Priming mitigated this reduction, although the highest value of the xylem potential was measured in the control treatment without saline stress (table 2).




Figure 1. Effect of priming on growth parameters of 40 days-old melon plants: leaf number (A), main stem length (B), leaf area (C), and total dry weight (D). [Control S0: non-stressed plants, irrigated with nutrient solution with 2.0 dS m⁻¹; Control S1: stressed plants, watered with 8.0 dS m⁻¹ nutrient solution; NaCl and CaCl₂: plants from primed seed during two days with NaCl and CaCl₂ respectively and watered with 8.0 dS m⁻¹ nutrient solution]. Different letters indicate a significant difference ($P \leq 0.05$) according to Tukey's test.




Figure 2. Melon plants of 20 days old. A: Control S0: non-stressed plants, irrigated with nutrient solution with 2.0 dS m⁻¹; B: Control S1: stressed plants, watered with 8.0 dS m⁻¹ nutrient solution; C and D: plants from primed seeds during two days with NaCl and CaCl₂ respectively and watered with 8.0 dS m⁻¹ nutrient solution.

Effect of priming on Na⁺/K⁺ Ratio and K⁺ leakage from roots

Na⁺/K⁺ ratio as well as K⁺ leakage by roots had a reduction in plants derived from primed seeds, compared with plants without previous treatment (Control S1). There were no significant differences between both salts (table 2). It was not possible to completely reverse the effect of salinity on these parameters.

Biochemical changes in plants

All biochemical parameters analyzed were affected by salinity (fig. 3). Priming allowed reversing partially this negative effect.

Reduction in total chlorophyll content, relative water content, and root viability were partially alleviated through priming treatments. In general, no differences were observed between the use of NaCl or CaCl₂ (fig. 3 A, B, and C).

An increase in relative electrolyte leakage and malondialdehyde (MDA) was measured in stressed plants. A lower increase was measured in plants derived from primed seeds (fig. 3 E and F).

Proline content, which is associated with the response to stress and with adaptation, was higher in stressed plants, and an even higher increase was registered in plants from primed seeds (fig. 3 D).

The activity of antioxidant enzymes increased under saline stress conditions and this increase was higher in priming treatments (fig. 3 G and H). The activity of peroxidase and catalase enzymes was higher in priming treatments. The NaCl treatment was even higher.

DISCUSSION

Priming conditions of the experiment were based on the research conducted by Nascimento (2003), who studied the effect of several treatments on melon germination at 17 and 25 °C. When comparing seed exposure to different osmotic potentials (Ψ_o , from -1.0 to -1.5 MPa) during different times (3, 6, 9, or 12 days), the best results were obtained with Ψ_o between -1.3 and -1.5 MPa for 3 days. On the other hand, Bradford (1986) suggested that Ψ_o lower than -2.0 MPa can damage seeds. Values close to 0 MPa could allow germination during treatment. In this work, an osmotic potential of -1.5 MPa and 2 days of priming allowed to reverse the negative effect of salt stress during the first vegetative stages of melon plants. With this osmotic potential, there was no damage to the seeds, and germination was prevented during the priming treatment.

The choice of the osmotic agents was based on previous work (Castañares and Bouzo, 2018) in which the effect of different agents (polyethylene glycol, KNO₃+K₃PO₄, NaCl and CaCl₂) on melon germination in saline stress was studied.

Germination reduction in saline conditions can be attributed to a combination of osmotic and toxic effects. Water absorption during imbibition is diminished with the reduction of Ψ_o caused by solutes (Ibrahim, 2016). On the other hand, high salinity can affect germination due to the toxic effects of Na⁺ and Cl⁻ on the viability of the embryo (Daszkowska-Golec, 2011).

Treatment	Ψ_x (atm)	Na ⁺ /K ⁺ ratio	K ⁺ leakage (mmol g ⁻¹ DW)
Control S0	-2.83 ± 2.29 c	0.022 ± 0.006 c	3.07 ± 0.33 c
Control S1	-8.50 ± 0.50 a	0.098 ± 0.004 a	35.85 ± 1.84 a
NaCl	-4.40 ± 0.36 b	0.056 ± 0.006 b	21.55 ± 1.06 b
CaCl ₂	-5.73 ± 0.25 ab	0.060 ± 0.004 b	24.91 ± 0.29 b

Table 2. Effect of priming on xylematic potential (Ψ_x), Na⁺/K⁺ ratio, and K⁺ leakage. [Control S0: non-stressed plants, irrigated with nutrient solution with 2.0 dS m⁻¹; Control S1: stressed plants, watered with 8.0 dS m⁻¹ nutrient solution; NaCl and CaCl₂: plants from primed seed during two days with NaCl and CaCl₂ respectively and watered with 8.0 dS m⁻¹ nutrient solution]. Different letters indicate a significant difference ($P \leq 0.05$) according to Tukey's test.




Figure 3. Effect of priming on total chlorophyll (A), relative water content (B) root viability (C), proline (D), relative electrolyte leakage (E), MDA content (F), peroxidase (G) and catalase (H) activity. [Control S0: non-stressed plants, irrigated with nutrient solution with 2.0 dS m⁻¹; Control S1: stressed plants, watered with 8.0 dS m⁻¹ nutrient solution; NaCl and CaCl₂: plants from primed seed during two days with NaCl and CaCl₂ respectively and watered with 8.0 dS m⁻¹ nutrient solution]. Different letters indicate a significant difference ($P \leq 0.05$) according to Tukey's test.

Germination improvement after priming (table 1) confirms the beneficial effect of this treatment. This effect can be attributed to the activation of metabolic processes that allow faster radicle emission while accelerating water absorption (Bewley and Black, 1994). Similar results were reported by Sivritepe *et al.* (1997) who studied melon germination in salinity conditions after priming with NaCl, registering an improvement in total germination and a reduction in the mean time of germination.

When comparing both priming durations (2 or 4 days) a decrease in germination was observed by extending the

imbibition time (table 1). Prolonged partial imbibition above the optimum for a given species will lead to a decrease of reserves, with the consequent reduction of vigor and viability of seeds, and irregular water absorption and loss of vital electrolytes for seeds (Jett *et al.*, 1996). Also, rapid seedling establishment might minimize crop risk due to environmental conditions or insect, and disease problems during field emergence.

From the analysis of the growth parameters (fig. 1) it is concluded that, although priming could not completely re-

verse the effect of salt stress, an increase in tolerance was recorded. Sivritepe *et al.* (2003) observed that with EC values of up to 9.0 dS m⁻¹, there was a difference in the dry weight of melon plants from primed seeds. At higher levels of salinity, there was no difference between treated and untreated plants.

Xylematic potential (Ψ_x) indicates water tension in plants' vessels which is related to the degree of stress (Selles and Ferreyra, 2005). The reduction of Ψ_x registered in plants growing under salinity conditions (table 2) confirms its effect on the hydric state of plants, due to the reduction of water availability (Shabala and Munns, 2012). The decrease in Ψ_x was lower in NaCl treatment, which suggests the activation of osmotic adjustment mechanisms, such as accumulation of compatible osmolytes, mainly proline (fig. 3 D), glycine betaine and sucrose, which reduce water potential of cells, protect against membrane damage and stabilize proteins and enzymes (Singh *et al.*, 2015). Proline accumulation in plants partially reduces the negative effects of stress given the ability of this molecule to participate in processes of osmotic adjustment and neutralization of ROS (Rejeb *et al.*, 2014). As observed in fig. 3 D, stressed plants increased proline content and an even higher content of this amino acid was measured in priming treatments. The increase in proline content in salt stressed plants was widely documented. In primed seeds, for example, was registered in pepper (*Capsicum annuum* L.) (Aloui *et al.*, 2014), rice (*Oryza sativa* L.) (Li and Zhang, 2012) and corn (*Zea mays* L.) (Bakht *et al.*, 2011).

Reduction of chlorophyll content in plants under saline stress can be related to a reduction of plant biomass and an increase in lipid peroxidation of the chloroplast's membrane due to oxidative stress, and the accumulation to toxic levels of ions that affect synthesis (Ashraf and Harris, 2013). The increase or reduction of chlorophyll content in saline stress has been proposed in melon as an indicator of tolerance or sensitivity to salinity, respectively (Romero *et al.*, 1997). There was a lower reduction of this pigment's content in primed treatments (fig. 3 A). When working with wheat (*Triticum aestivum* L.), Farooq *et al.* (2013), found a high correlation between the reduction of chlorophyll content in plants under salt stress and the degree of lipid peroxidation and membrane damage. Our results confirm this relationship (fig. 3 A, E, and F).

Reduction of relative water content (RWC) in stressed plants (fig. 3 B) can be related to lower water absorption due to damage of the root system or the low Ψ of the substrate (Kukreja *et al.*, 2005). In this way, the increase of RWC in priming treatments can be explained, on the one hand, by the lower root damage, evidenced with the highest values of reaction to TTC (fig. 3 C), as an indicator of root viability (Ruf and Brunner, 2003).

Stress in plants triggers numerous processes that lead to the generation of ROS. These ROS do not necessarily have a negative effect on plants, but they do have an important function in intracellular communication that allows the best acclimatization to environmental stress (Rejeb *et*

al., 2014). However, excessive accumulation causes the initiation of a significant number of auto oxidative chain reactions, mainly lipid peroxidation, DNA damage and protein degradation (Mittler, 2002), since these are molecules with a highly unstable configuration. Therefore, radicals quickly react with other molecules, generating more free radicals (Ashraf, 2009). Hydrogen peroxide (H₂O₂), which was not measured here, is the most harmful ROS and has been widely studied because of its relative stability (Mhamdi *et al.*, 2010). Antioxidant enzymes, particularly peroxidases (POX) and catalases (CAT) are very efficient at degrading H₂O₂ (Li *et al.*, 2014). Therefore, the high values measured here of these enzymes (fig. 3 G and H) indicate effective mitigation of the possible damage of H₂O₂ on melon plants. Saline stress caused an increase in the activity of POX (fig. 3 G) and CAT (fig. 3 H) enzymes. Priming led to an even greater increase, primarily for NaCl followed by CaCl₂. These results agree with was reported by Islam *et al.* (2015) who studied the response of priming with CaCl₂ and KCl in two wheat cultivars (*Triticum aestivum* L.) grown in saline soil, and reported a higher antioxidant activity. Salah *et al.* (2015) cultivated rice plants (*Oryza sativa* L.) under toxic levels of Zn, and observed a higher POX and CAT activity, and lower levels of MDA, in plants from seeds primed with PEG.

Under high salinity conditions, Na⁺ has the ability to compete with K⁺. Many physiological studies have demonstrated that Na⁺ toxicity is caused by the ability of Na⁺ to compete for K⁺ binding sites to disrupt K⁺ homeostasis (Hasegawa, 2013). One of the main characteristics of salt tolerant plants is the ability to maintain a low Na⁺/K⁺ ratio (Tester and Davenport, 2003). In our experiments, plant growing in saline conditions increased Na⁺/K⁺ (table 2). This can be explained by a greater Na⁺ accumulation in tissues and an increase in K⁺ leakage by cells (Poustini and Siosemardeh, 2004). The reduction of Na⁺/K⁺ ratio in plants from primed seeds contributes to a better adaptation to salinity. In wheat, Abbasdokht and Edalatpisheh (2012) observed that plants from primed seeds had low Na⁺/K⁺ ratio, while the untreated plants exhibited higher ratios. Working with tomato (*Solanum lycopersicum* L.), Cayuela *et al.* (1996) reported that priming with NaCl allowed reducing Na⁺ content in plants that best grow in salinity. Plants growing in salt stress can experience an irreversible loss of K⁺, which is particularly important in roots. This is related to the loss of integrity of membranes, reducing the ability to retain K⁺ (Demidchik *et al.*, 2014). Low K⁺ leakage from roots is accepted as an indicator of salt tolerance (Chen *et al.*, 2005). The lower K⁺ release measured in priming treatments (table 2) confirms the greater stability of membranes, also estimated from relative electrolyte leakage (fig. 3 E) and MDA (fig. 3 F). K⁺ is recognized as an important factor limiting yield and quality of crops, and functions include maintenance of cell turgor, growth, activation of numerous enzymes, stabilization of protein synthesis, the formation of membrane potential, and maintenance of cytosolic pH (Cakmak, 2005). Relative electrolyte leakage is considered an estimator of membrane integrity (Demidchik *et al.*, 2014). A greater release of electrolytes, estimated from EC, indicates membrane damage. This was measured in stressed plants (Control S1)

while priming allowed reducing this damage (fig. 3 E). The final product of polyunsaturated fatty acids oxidation is malondialdehyde (MDA). This molecule reacts with thiobarbituric acid (TBA) and generated a pink compound detectable by spectrophotometry (Hodges *et al.*, 1999). The increase of the MDA level in the conditions of the experiment (fig. 3 F) confirms the membrane damage, which corroborates the results discussed above by the electrolyte leakage measurement (fig. 3 E). Thus, priming allowed improving the integrity of membranes under the conditions of the experiment. This response is in agreement with what was observed by Randhir and Shetty (2005) in corn (*Zea mays* L.) and Amooaghaie (2011) in alfalfa (*Medicago sativa* L.).

CONCLUSIONS

The results of this study show that priming induces biochemical changes in melon seedlings increasing the adaptive response to salinity. Melon is cultivated intensively and with complementary irrigation, which promotes the accumulation of salts in the soil. Given that the studied salts have a relatively low cost and are easy to manipulate, the information obtained could be of interest to producers.

REFERENCES

- ABBASDOKHT, H.; EDALATPISHEH, M.R. 2012. The effect of priming and salinity on physiological and chemical characteristics of wheat (*Triticum aestivum* L.). *Desert* 17, 183–192.
- AEBI, H. 1984. Catalase in vitro, in: PACKER, L. (Ed.). *Methods in Enzymology*. Academic Press, New York. 121–126 pp.
- ALOUI, H.; SOUGUIR, M.; LATIQUE, S.; HANNACHI, C. 2014. Germination and growth in control and primed seeds of pepper as affected by salt stress. *Cercet. Agron. Mold.* 47, 83–95.
- AMOOAGHAIE, R. 2011. The effect of hydro and osmopriming on alfalfa seed germination and antioxidant defenses under salt stress. *African J. Biotechnol.* 10, 6269–6275.
- ASHRAF, M. 2009. Biotechnological approach of improving plant salt tolerance using antioxidants as markers. *Biotechnol. Adv.* 27, 84–93. <https://doi.org/10.1016/j.biotechadv.2008.09.003>
- ASHRAF, M.; HARRIS, P.J.C. 2013. Photosynthesis under stressful environments: an overview. *Photosynthetica* 51, 163–190.
- BAKHT, J.; SHAFI, M.; JAMAL, Y.; SHER, H. 2011. Response of maize (*Zea mays* L.) to seed priming with NaCl and salinity stress. *Spanish J. Agric. Res.* 9, 252–261.
- BATES, L.S.; WALDREN, R.P.; TEARE, I.D. 1973. Rapid determination of free proline for water-stress studies. *Plant Soil* 39, 205–207. <https://doi.org/10.1007/BF00018060>
- BEWLEY, J.D.; BLACK, M. 1994. *Seeds*, in: *Seeds*. Springer, New York. 1–33 pp.
- BOKOCHOY-NBOK, H.; BOKOCHOY, A. 1991. Response of melon plants to salt: 1. Growth, morphology and root membrane properties. *J. Plant Physiol* 139, 100–105.
- BOTÍA, P.; CARVAJAL, M.; CERDÁ, A.; MARTÍNEZ, V. 1998. Response of eight *Cucumis melo* cultivars to salinity during germination and early vegetative growth. *Agronomie* 18, 503–513. <https://doi.org/10.1051/agro:19980801>
- BRADFORD, K.J. 1986. Manipulation of seed water relations via osmotic priming to improve germination under stress conditions. *HortScience (USA)* 21, 1105–1113.
- CAKMAK, I. 2005. The role of potassium in alleviating detrimental effects of abiotic stresses in plants. *J. Plant Nutr. Soil Sci.* 168, 521–530.
- CAMPBELL, C.R.; PLANK, C.O. 1998. Preparation of plant tissue for laboratory analysis, in: KALRA, Y.P. (Ed.), *Handbook of Methods for Plant Analysis*. Taylor & Francis Group, New York. 51–63 pp.
- CAO, Y.Y.; YANG, M.T.; CHEN, S.Y.; ZHOU, Z.Q.; LI, X.; WANG, X.J.; BAI, J.G. 2015. Exogenous sucrose influences antioxidant enzyme activities and reduces lipid peroxidation in water-stressed cucumber leaves. *Biol. Plant.* 59, 147–153.
- CARVAJAL, M.; DEL AMOR, F.M.; FERNANDEZ-BALLESTER, G.; MARTÍNEZ, V.; CERDÁ, A. 1998. Time course of solute accumulation and water relations in muskmelon plants exposed to salt during different growth stages. *Plant Sci.* 138, 103–112.
- CASTAÑARES, J.L.; BOUZO, C.A. 2018. Effect of different priming treatments and priming durations on melon germination behavior under suboptimal conditions. *Open Agric.* 3, 386–392.
- CASTAÑARES, J.L.; BOUZO, C.A. 2017. Effect of priming on germination and initial growth of melon plants under salt stress. *FAME Sección Ciencias Agrar.* 16, 7–16.
- CAYUELA, E.; PÉREZ-ALFOCEA, F.; CARO, M.; BOLARÍN, M.C. 1996. Priming of seeds with NaCl induces physiological changes in tomato plants grown under salt stress. *Physiol. Plant.* 96, 231–236. <https://doi.org/10.1111/j.1399-3054.1996.tb00207.x>
- CHEN, Z.; NEWMAN, I.; ZHOU, M.; MENDHAM, N.; ZHANG, G.; SHABALA, S. 2005. Screening plants for salt tolerance by measuring K⁺ flux: a case study for barley. *Plant. Cell Environ.* 28, 1230–1246.
- CLEMENSSON-LINDELL, A. 1994. Triphenyltetrazolium chloride as an indicator of fine-root vitality and environmental stress in coniferous forest stands: applications and limitations. *Plant Soil* 159, 297–300.
- DASZKOWSKA-GOLEC, A. 2011. Arabidopsis seed germination under abiotic stress as a concert of action of phytohormones. *Omi. a J. Integr. Biol.* 15, 763–774.
- DEMIDCHIK, V.; STRALTISOVA, D.; MEDVEDEV, S.S.; POZHVANOV, G.A.; SOKOLIK, A.; YURIN, V. 2014. Stress-induced electrolyte leakage: the role of K⁺-permeable channels and involvement in programmed cell death and metabolic adjustment. *J. Exp. Bot.* 65, 1259–1270.
- DI RIENZO, J.A.; CASANOVES, F.; BALZARINI, M.G.; GONZÁLEZ, L.; TABLADA, M.; ROBLEDO, Y.C.W. 2011. InfoStat versión 2011. Grup. InfoStat, FCA, Univ. Nac. Córdoba, Argentina. (Available at: <http://www.infostat.com.ar> verified: December 2019) 195–199.
- DIONISIO-SESE, M.L.; TOBITA, S. 1998. Antioxidant responses of rice seedlings to salinity stress. *Plant Sci.* 135, 1–9.
- FARHOUDI, R.; SHARIFZADEH, F. 2006. The effects of NaCl priming on salt tolerance in canola (*Brassica napus* L.) seedlings grown under saline conditions. *Indian J Crop Sci* 1, 74–78.
- FAROOQ, M.; IRFAN, M.; AZIZ, T.; AHMAD, I.; CHEEMA, S.A. 2013. Seed priming with ascorbic acid improves drought resistance of wheat. *J. Agron. Crop Sci.* 66, 12–22.
- FRANCO, J.A.; ESTEBAN, C.; RODRIGUEZ, C. 1993. Effects of salinity on various growth stages of muskmelon cv. Revigal. *J. Hortic. Sci.* 68, 899–904.
- GEORGE, P. 1953. The chemical nature of the second hydrogen peroxide compound formed by cytochrome c peroxidase and horseradish peroxidase. 1. Titration with reducing agents. *Biochem. J.* 54, 267–276.
- HASEGAWA, P.M. 2013. Sodium (Na⁺) homeostasis and salt tolerance of plants. *Environ. Exp. Bot.* 92, 19–31.

- HEATH, R.L.; PACKER, L. 1968. Photoperoxidation in isolated chloroplasts: I. Kinetics and stoichiometry of fatty acid peroxidation. *Arch. Biochem. Biophys.* 125, 189–198.
- HOAGLAND, D.R.; ARNON, D.I. 1950. The Water-Culture Method for Growing Plants without Soil. *Calif. Agric. Exp. Stn.* 51, 914–916.
- HODGES, D.M.; DELONG, J.M.; FORNEY, C.F.; PRANGE, R.K. 1999. Improving the thiobarbituric acid-reactive-substances assay for estimating lipid peroxidation in plant tissues containing anthocyanin and other interfering compounds. *Planta* 207, 604–611.
- HUANG, C.H.; ZONG, L.; BUONANNO, M.; XUE, X.; WANG, T.; TEDESCHI, A. 2012. Impact of saline water irrigation on yield and quality of melon (*Cucumis melo* cv. Huanghem) in northwest China. *Eur. J. Agron.* 43, 68–76.
- IBRAHIM, E.A. 2016. Seed priming to alleviate salinity stress in germinating seeds. *J. Plant Physiol.* 192, 38–46. <https://doi.org/10.1016/j.jplph.2015.12.011>
- IQBAL, M.; UL HAQ, Z.; JAMIL, Y.; NISAR, J. 2016. Pre-sowing seed magnetic field treatment influence on germination, seedling growth and enzymatic activities of melon (*Cucumis melo* L.). *Bio-catal. Agric. Biotechnol.* 6, 176–183.
- ISLAM, F.; YASMEEN, T.; ALI, S.; ALI, B.; FAROOQ, M.A.; GILL, R.A. 2015. Priming-induced antioxidative responses in two wheat cultivars under saline stress. *Acta Physiol. Plant.* 37, 153–165.
- ISTA. 2013. International rules for seed testing. *Seed Sci. Technol.* 27, 25–30.
- JETT, L.W.; WELBAUM, G.E.; MORSE, R.D. 1996. Effects of matric and osmotic priming treatments on broccoli seed germination. *J. Am. Soc. Hortic. Sci.* 121, 423–429.
- JISHA, K.C.; VIJAYAKUMARI, K.; PUTHUR, J.T. 2013. Seed priming for abiotic stress tolerance: an overview. *Acta Physiol. Plant.* 35, 1381–1396.
- KUKREJA, S.; NANDWAL, A.S.; KUMAR, N.; SHARMA, S.K.; UNVI, V.; SHARMA, P.K. 2005. Plant water status, H_2O_2 scavenging enzymes, ethylene evolution and membrane integrity of *Cicer arietinum* roots as affected by salinity. *Biol. Plant.* 49, 305–308.
- LI, C.; TAN, D.X.; LIANG, D.; CHANG, C.; JIA, D.; MA, F. 2014. Melatonin mediates the regulation of ABA metabolism, free-radical scavenging, and stomatal behaviour in two *Malus* species under drought stress. *J. Exp. Bot.* 66, 669–680.
- LI, X.; ZHANG, L. 2012. SA and PEG-induced priming for water stress tolerance in rice seedling, in: *Information Technology and Agricultural Engineering*. Springer, 881–887 pp.
- LICHTENTHALER, H.K.; WELLBURN, A.R. 1983. Determinations of total carotenoids and chlorophylls a and b of leaf extracts in different solvents. *Biochem. Soc. Trans.* 11, 591–592.
- MHAMDI, A.; QUEVAL, G.; CHAOUCH, S.; VANDERAUWERA, S.; VAN BREUSEGEM, F.; NOCTOR, G. 2010. Catalase function in plants: a focus on Arabidopsis mutants as stress-mimic models. *J. Exp. Bot.* 61, 4197–4220.
- MITTLER, R. 2002. Oxidative stress, antioxidants and stress tolerance. *Trends Plant Sci.* 7, 405–410.
- MUNNS, R. 2005. Genes and salt tolerance: bringing them together. *New Phytol.* 167, 645–663.
- NASCIMENTO, W.M. 2003. Muskmelon seed germination and seedling development in response to seed priming. *Sci. Agric.* 60, 71–75.
- NUKAYA, A.; MASUI, M.; ISHIDA, A. 1984. Salt tolerance of muskmelons as affected by diluted sea water applied at different growth stages in nutrient solution culture. *J. Japanese Soc. Hortic. Sci.* 53, 168–175.
- PAPARELLA, S.; ARAÚJO, S.S.; ROSSI, G.; WIJAYASINGHE, M.; CARBONERA, D.; BALESTRAZZI, A. 2015. Seed priming: state of the art and new perspectives. *Plant Cell Rep.* 34, 1281–1293. <https://doi.org/10.1007/s00299-015-1784-y>
- PASTOR, V.; BALMER, A.; GAMIR, J.; FLORS, V.; MAUCH-MANI, B. 2014. Preparing to fight back: generation and storage of priming compounds. *Front. Plant Sci.* 5, 1–12.
- POUSTINI, K.; SIOSEMARDEH, A. 2004. Ion distribution in wheat cultivars in response to salinity stress. *F. rops Res* 85, 125–133.
- QADIR, M.; QUILLÉROU, E.; NANGIA, V.; MURTAZA, G.; SINGH, M.; THOMAS, R.J.; DRECHSEL, P.; NOBLE, A.D. 2014. Economics of salt induced land degradation and restoration, in: *Natural Resources Forum*. Wiley Online Library. 282–295 pp.
- RANDHIR, R.; SHETTY, K. 2005. Developmental stimulation of total phenolics and related antioxidant activity in light-and dark-germinated corn by natural elicitors. *Process Biochem.* 40, 1721–1732.
- RECHE MÁRMOL, J. 2008. Cultivo del melón en invernadero, Generalitat Valenciana Conselleria d'Agricultura i Pesca. Valencia, España. Secretaría General Técnica. Servicio de Publicaciones y Divulgación, Andalucía.
- REJEB, K.B.; ABDELLY, C.; SAVOURÉ, A. 2014. How reactive oxygen species and proline face stress together. *Plant Physiol. Biochem.* 80, 278–284.
- ROMERO, L.; BELAKBIR, A.; RAGALA, L.; RUIZ, J.M. 1997. Response of plant yield and leaf pigments to saline conditions: Effectiveness of different rootstocks in melon plants (*Cucumis melo* L.). *Soil Sci. Plant Nutr.* 43, 855–862. <https://doi.org/10.1080/00380768.1997.10414652>
- RUF, M.; BRUNNER, I. 2003. Vitality of tree fine roots: reevaluation of the tetrazolium test. *Tree Physiol.* 23, 257–263.
- SALAH, S.M.; YAJING, G.; DONGDONG, C.; JIE, L.; AAMIR, N.; QIJUAN, H.; WEIMIN, H.; MINGYU, N.; JIN, H. 2015. Seed priming with polyethylene glycol regulating the physiological and molecular mechanism in rice (*Oryza sativa* L.) under nano-ZnO stress. *Sci. Rep.* 5, 1–14.
- SCHNEIDER, C.A.; RASBAND, W.S.; ELICEIRI, K.W. 2012. NIH Image to ImageJ: 25 years of image analysis. *Nat. Methods* 9, 671–675.
- SCHOLANDER, P.F.; HAMMEL, H.T.; HEMMINGSEN, E.A.; BRADSTREET, E.D. 1964. Hydrostatic pressure and osmotic potential in leaves of mangroves and some other plants. *Proc. Natl. Acad. Sci.* 52, 119–125.
- SELLES, G.; FERREYRA, R. 2005. Criterios para controlar el riego en uva de mesa. INIA (ed). *Manejo Riego y Suelo en Vides para Vino y Mesa*. INIA, Santiago, Chile 15–53.
- SHABALA, S.; MUNNS, R. 2012. Salinity stress: physiological constraints and adaptive mechanism, in: SHABALA, S. (Ed.). *Plant Stress Physiology*. CAB International, London. 59–93 pp.
- SINGH, M.; KUMAR, J.; SINGH, S.; SINGH, V.P.; PRASAD, S.M. 2015. Roles of osmoprotectants in improving salinity and drought tolerance in plants: a review. *Rev. Environ. Sci. Bio/Technology* 14, 407–426.
- SIVRITEPE, H.Ö.; SIVRITEPE, N.; ERIŞ, A.; TURHAN, E. 2005. The effects of NaCl pre-treatments on salt tolerance of melons grown under long-term salinity. *Sci. Hortic. (Amsterdam)*. 106, 568–581.
- SIVRITEPE, N.; SIVRITEPE, H.Ö.; ERIS, A. 2003. The effects of NaCl priming on salt tolerance in melon seedlings grown under saline conditions. *Sci. Hortic. (Amsterdam)*. 97, 229–237.
- SIVRITEPE, Ö.H.; ERIS, A.; SIVRITEPE, N. 1997. The effects of priming treatments on salt tolerance in melon seeds, in: *I International Symposium on Cucurbits* 492. 287–296 pp.
- TANVEER, A.; SALMAN ARSHAD, M.; AYUB, M.; MANSOOR JAVOID, M.; YASEEN, M. 2012. Effect of temperature, light, sali-

nity, drought stress and seeding depth on germination of *Cucumis melo* var. *agrestis*. Pakistan J. Weed Sci. Res. 18, 445–459.

TEDESCHI, A.; LAVINI, A.; RICCARDI, M.; PULVENTO, C.; D'ANDRIA, R. 2011. Melon crops (*Cucumis melo* L., cv. Tendral) grown in a mediterranean environment under saline-sodic conditions: Part I. Yield and quality. Agric. water Manag. 98, 1329–1338.

TESTER, M.; DAVENPORT, R. 2003. Na⁺ tolerance and Na⁺ transport in higher plants. Ann. Bot. 91, 503–527.

WOJTYLA, Ł.; LECHOWSKA, K.; KUBALA, S.; GARNCZARSKA, M. 2016. Molecular processes induced in primed seeds—increasing the potential to stabilize crop yields under drought conditions. J. Plant Physiol. 203, 116–126.

YADAV, S.; IRFAN, M.D.; AHMAD, A.; HAYAT, S. 2011. Causes of salinity and plant manifestations to salt stress: A review. J. Environ. Biol. 32, 667–685.

YASAR, F.; KUSVURAN, S.; ELLIALTIOGLU, S. 2006. Determination of anti-oxidant activities in some melon (*Cucumis melo* L.) varieties and cultivars under salt stress. J. Hortic. Sci. Biotechnol. 81, 627–630.

ZHENG, M.; TAO, Y.; HUSSAIN, S.; JIANG, Q.; PENG, S.; HUANG, J.; CUI, K.; NIE, L. 2016. Seed priming in dry direct-seeded rice: consequences for emergence, seedling growth and associated metabolic events under drought stress. Plant Growth Regul. 78, 167–178.